

# Programming Languages and Compilers (CS 421)

Elsa L Gunter  
 2112 SC, UIUC  
<http://www.cs.uiuc.edu/class/fa06/cs421/>

Based in part on slides by Mattox Beckman, as updated by Vikram Adve and Gul Agha

## Type Inference - The Problem

- Given an expression  $e$ , and a typing environment  $\Gamma$ , does there exist a type  $\tau$  such that the judgment
 
$$\Gamma \vdash e : \tau$$
 is valid - ie., follows from the typing rules?

Elsa L. Gunter

## Type Inference - Outline

- Begin by assigning a type variable as the type of the whole expression
- Decompose the expression into component expressions
- Use typing rules to generate constraints on components and whole
- Recursively gather additional constraints to guarantee a solution for components
- Solve system of constraints to generate a substitution
- Apply substitution to orig. type var. to get answer

Elsa L. Gunter

## Type Inference - Example

- What type can we give to
 
$$\text{fun } x \rightarrow \text{fun } f \rightarrow f \ x?$$
- Start with a type variable and then look at the way the term is constructed

Elsa L. Gunter

## Type Inference - Example

- First approximate:
 
$$[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f \ x) : \alpha$$
- Second approximate: use fun rule
 
$$\frac{[x : \beta] \vdash (\text{fun } f \rightarrow f \ x) : \gamma}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f \ x) : \alpha}$$
- $\alpha \equiv (\beta \rightarrow \gamma)$

Elsa L. Gunter

## Type Inference - Example

- Third approximate: use fun rule
 
$$\frac{\frac{[f : \delta ; x : \beta] \vdash (f \ x) : \varepsilon}{[x : \beta] \vdash (\text{fun } f \rightarrow f \ x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f \ x) : \alpha}$$
- $\alpha \equiv (\beta \rightarrow \gamma); \gamma \equiv (\delta \rightarrow \varepsilon)$

Elsa L. Gunter

## Type Inference - Example

- Fourth approximate: use app rule

$$\frac{\frac{\frac{[f : \delta; x : \beta] \vdash f : \varphi \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \varphi}{[f : \delta; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv (\delta \rightarrow \varepsilon)$

Elsa L. Gunter

## Type Inference - Example

- Fifth approximate: use var rule

$$\frac{\frac{\frac{[f : \delta; x : \beta] \vdash f : \varphi \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \varphi}{[f : \delta; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv (\delta \rightarrow \varepsilon)$ ;  $\delta \equiv (\varphi \rightarrow \varepsilon)$

Elsa L. Gunter

## Type Inference - Example

- Sixth approximate: use var rule

$$\frac{\frac{\frac{\frac{[f : \delta; x : \beta] \vdash f : \varphi \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \varphi}{[f : \delta; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv (\delta \rightarrow \varepsilon)$ ;  $\delta \equiv (\varphi \rightarrow \varepsilon)$ ;  $\varphi \equiv \beta$

Elsa L. Gunter

## Type Inference - Example

- Done building proof tree; now solve!

$$\frac{\frac{\frac{\frac{[f : \delta; x : \beta] \vdash f : \varphi \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \varphi}{[f : \delta; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv (\delta \rightarrow \varepsilon)$ ;  $\delta \equiv (\varphi \rightarrow \varepsilon)$ ;  $\varphi \equiv \beta$

Elsa L. Gunter

## Type Inference - Example

- Type unification; solve like linear equations;

$$\frac{\frac{\frac{\frac{[f : \delta; x : \beta] \vdash f : \varphi \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \varphi}{[f : \delta; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv (\delta \rightarrow \varepsilon)$ ;  $\delta \equiv (\varphi \rightarrow \varepsilon)$ ;  $\varphi \equiv \beta$

Elsa L. Gunter

## Type Inference - Example

- Eliminate  $\varphi$ :

$$\frac{\frac{\frac{\frac{[f : \delta; x : \beta] \vdash f : \beta \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \beta}{[f : \delta; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv (\delta \rightarrow \varepsilon)$ ;  $\delta \equiv (\beta \rightarrow \varepsilon)$ ;  $\varphi \equiv \beta$

Elsa L. Gunter

## Type Inference - Example

- Next eliminate  $\delta$  :

$$\frac{\frac{\frac{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash f : \beta \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \beta}{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : \gamma}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow \gamma)$ ;  $\gamma \equiv ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon)$ ;  $\delta \equiv (\beta \rightarrow \varepsilon)$  ;

Elsa L. Gunter

## Type Inference - Example

- Next eliminate  $\gamma$  :

$$\frac{\frac{\frac{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash f : \beta \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \beta}{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon)}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : \alpha}}$$

- $\alpha \equiv (\beta \rightarrow ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon))$ ;  $\gamma \equiv ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon)$ ;

Elsa L. Gunter

## Type Inference - Example

- Next eliminate  $\alpha$  :

$$\frac{\frac{\frac{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash f : \beta \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \beta}{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon)}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : (\beta \rightarrow ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon))}}$$

- $\alpha \equiv (\beta \rightarrow ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon))$ ;

Elsa L. Gunter

## Type Inference - Example

- No more equations to solve; we are done

$$\frac{\frac{\frac{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash f : \beta \rightarrow \varepsilon \quad [f : \delta; x : \beta] \vdash x : \beta}{[f : \beta \rightarrow \varepsilon; x : \beta] \vdash (f x) : \varepsilon}}{[x : \beta] \vdash (\text{fun } f \rightarrow f x) : ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon)}}{[] \vdash (\text{fun } x \rightarrow \text{fun } f \rightarrow f x) : (\beta \rightarrow ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon))}}$$

- Any instance of  $(\beta \rightarrow ((\beta \rightarrow \varepsilon) \rightarrow \varepsilon))$  is a valid type

Elsa L. Gunter

## Type Inference - The Problem

- Given an expression  $e$ , and a typing environment  $\Gamma$ , does there exist a type  $\tau$  such that the judgment

$$\Gamma \vdash e : \tau$$

is valid - ie., follows from the typing rules?

Elsa L. Gunter

## Type Inference Algorithm

Let  $\text{has\_type}(\Gamma, e, \tau) = S$

- $\Gamma$  is a typing environment
- $e$  is an expression
- $\tau$  is a (generalized) type,
- $S$  is a set of equations between generalized types

Elsa L. Gunter

## Type Inference Algorithm

- Let  $\text{has\_type}(\Gamma, e, \tau) = S$ 
  - $\Gamma$  is a typing environment,
  - $e$  is an expression,
  - $\tau$  is a (generalized) type,
  - $S$  is a set of equations between generalized types.  
**Idea:**  $S$  is the constraints on type variables necessary for  $\Gamma \vdash e : \tau$
- Let  $\text{Unif}(S)$  be a substitution of generalized types for type variables solving  $S$
- Solution:  $\text{Unif}(S)(\Gamma) \vdash e : \text{Unif}(S)(\tau)$

Elsa L. Gunter

## Type Inference Algorithm

$\text{has\_type}(\Gamma, \text{exp}, \tau) =$

- Case  $\text{exp}$  of
  - Var  $v \rightarrow$  return  $\{\tau \equiv \Gamma(v)\}$
  - Const  $c \rightarrow$  return  $\{\tau \equiv \sigma\}$  where  $\Gamma \vdash c : \sigma$  by the constant rules
  - fun  $x \rightarrow e \rightarrow$ 
    - Let  $\alpha, \beta$  be fresh variables
    - Let  $S = \text{has\_type}([x: \alpha] \cup \Gamma, e, \beta)$
    - Return  $\{\tau \equiv \alpha \rightarrow \beta\} \cup S$

Elsa L. Gunter

## Type Inference Algorithm (cont)

- Case  $\text{exp}$  of
  - App  $(e_1 e_2) \rightarrow$ 
    - Let  $\alpha$  be a fresh variable
    - Let  $S_1 = \text{has\_type}(\Gamma, e_1, \alpha \rightarrow \tau)$
    - Let  $S_2 = \text{has\_type}(\Gamma, e_2, \alpha)$
    - Return  $S_1 \cup S_2$

Elsa L. Gunter

## Type Inference Algorithm (cont)

- Case  $\text{exp}$  of
  - let  $x = e_1$  in  $e_2 \rightarrow$ 
    - Let  $\alpha$  be a fresh variable
    - Let  $S_1 = \text{has\_type}(\Gamma, e_1, \alpha)$
    - Let  $S_2 =$   
 $\text{has\_type}([x: \alpha] \cup \Gamma, e_2, \tau)$
    - Return  $S_1 \cup S_2$

Elsa L. Gunter

## Type Inference Algorithm (cont)

- Case  $\text{exp}$  of
  - let rec  $x = e_1$  in  $e_2 \rightarrow$ 
    - Let  $\alpha$  be a fresh variable
    - Let  $S_1 = \text{has\_type}([x: \alpha] \cup \Gamma, e_1, \alpha)$
    - Let  $S_2 = \text{has\_type}([x: \alpha] \cup \Gamma, e_2, \tau)$
    - Return  $S_1 \cup S_2$

Elsa L. Gunter

## Type Inference Algorithm (cont)

- To infer a type, introduce  $\text{type\_of}$
- Let  $\alpha$  be a fresh variable
- $\text{type\_of}(\Gamma, e) =$ 
  - Let  $S = \text{has\_type}(\Gamma, e, \alpha)$
  - Return  $\text{Unif}(S)(\alpha)$
- Need an algorithm for Unif

Elsa L. Gunter